

On the Ordering Instability of Weakly-Interacting Electrons in a Dirty Metal

Chetan Nayak and Xiao Yang

Department of Physics and Astronomy, University of California at Los Angeles, Los Angeles, CA 90095-1547

(Dated: February 2, 2008)

In a dirty metal, electron-electron interactions in the spin-triplet channel lead to singular corrections to a variety of physical quantities. We show that these singularities herald the emergence of ferromagnetism. We calculate the effective action for the magnetic moment of weakly-interacting electrons in a dirty metal and show that a state with finite ferromagnetic moment minimizes this effective action. The saddle-point approximation is exact in an appropriate large- N limit. We discuss the physics of the ferromagnetic state with particular regard to thermal fluctuations and localization effects.

PACS numbers: 71.10.Hf, 71.10.-w, 71.30.+h, 73.23.-b

I. INTRODUCTION.

The presence in a solid of randomly-distributed imperfections causes its electrons to propagate diffusively rather than ballistically. As a result, when two electrons interact, they tarry in each other's vicinity, and become more strongly correlated¹. As shown by Finkelstein² and Castellani, *et al.*^{3,4}, this effect leads to an instability of the diffusive Fermi liquid fixed point (see also refs. 5,6,7) which manifests itself through divergent corrections to a variety of physical quantities. The instability exists for arbitrarily weak coupling and for arbitrarily weak disorder. Its simplest incarnation is in an electron system in a weak magnetic field $\mathbf{B}$ which suppresses quantum interference effects and relegates them to subleading order in the dimensionless resistance⁸ t (measured in units of h/e^2) while interactions make a contribution at leading order. Since the relevant interaction is in the spin-triplet particle-hole channel, the g factor must be tuned to zero if $\mathbf{B} \neq 0$ so that there is no Zeeman splitting, which would cut off the divergences. Though this instability was identified nearly 20 years ago, there is, at present, no consensus on its outcome.

The recent discovery of evidence for a metal-insulator transition in two-dimensional electron gases (2DEGs) in Si-MOSFETs^{9,10}, GaAs¹¹ heterostructures, and Si-Ge¹² has led to a re-examination of the problem of interacting electrons in a random potential. The interesting phenomena seen in these experiments occur in a regime in which interactions are strong (large r_s) and quantum interference effects are likely to be important because the conductivity is on the order of the quantum limit $\sim e^2/h$ and $B = 0$. Though a number of interesting theories^{13,14,15,16,17,18} have been proposed, none have provided a complete explanation of the experimental data, which isn't surprising, given that the presumably simpler problem of weakly-interacting electrons in a weak random potential with $\mathbf{B} \neq 0$, $g = 0$ remains unsolved. It is this simpler problem which is the subject of this paper.

Though it is unresolved, the instability caused by the triplet interaction is, in fact, a crime with only one plausible suspect: ferromagnetism. Since the diffusive Fermi

liquid is unstable, the system must settle into some other phase at low temperature; since the instability is in the spin-triplet channel, it is presumably one in which the spins are ordered. The basic interaction in this problem is exchange, which is ferromagnetic. There is no available mechanism which generates antiferromagnetic interactions. To see this, note that the instability occurs when the conductance is large, while superexchange occurs in or near an insulating state (since it arises when the kinetic energy is treated as a perturbation). There would be RKKY interactions between localized spins, such as magnetic impurities, if we had any, but we don't. In the absence of any antiferromagnetic interactions, we rule out spin density-waves (SDWs)¹⁹, spin glass order, and random singlet phases²⁰. The latter two furthermore require large effective disorder while the instability of interest here is manifest for small effective disorder²¹.

It has been suggested that the instability of the triplet interaction signals the formation of 'local moments'⁴. In the weak-interaction, weak-disorder limit, however, the instability of the diffusive Fermi liquid only develops at large length scales, so the resulting magnetic moments are anything but 'local'. Local moments might be expected when the energy cost for flipping a spin at short length scales is very small. This might occur near the Stoner transition of the clean system²², where the critical susceptibility, $\chi \sim 1/T^\gamma$, is of Curie form for a mean-field exponent $\gamma = 1$. There is also a substantial local moment regime with Curie susceptibility in the large- U Hubbard model (because of the weakness of superexchange for large U) and doped semiconductors (because of competing ferromagnetic and antiferromagnetic interactions). However, these scenarios are not relevant to the presumably simpler, though still non-trivial, problem of the weakly-interacting, weakly-disordered limit.

This leaves us with one remaining possibility and, indeed, a number of authors have proposed that the singularity of the triplet interaction channel leads to ferromagnetism. The basic underlying physics is simply that diffusing electrons interact more strongly because they move more slowly and, as a result, the interaction energy gained by (at least partially) polarizing the system *always* outweighs the single-particle energy cost, in contrast to

a clean system, where the interaction energy gain is a constant U which must exceed the single-particle energy cost $1/N_F$, i.e. the Stoner criterion (N_F is the density of states at the Fermi energy). Andreev and Kameney²³ proposed such a mechanism for finite-size systems by computing the disorder enhancement of the interaction matrix element for a state with two electrons in single-particle states separated by small energies. Heuristically, one can say that the two wavefunctions are concentrated in the same minima of the potential. Chamon and Mucciolo²⁴ found a ferromagnetic saddle-point solution of the one-loop effective action of Finkelstein's theory. Their calculation generalizes Stoner's by including quantum corrections resulting from interactions between diffusion modes. Belitz and Kirkpatrick²⁵ advanced the same hypothesis when they noted that the phase transition which they expected to be driven by the growth of the triplet interaction is of the same form as a transition to ferromagnetism in a disordered system.

In this paper, we calculate the effective action for the magnetization in a weakly-interacting electron system in a slightly dirty metal. The effective action is calculated to lowest order in the dimensionless resistance t . It exhibits the physics described above and is clearly unstable to the formation of a ferromagnetic moment. In principle, the emergence of some other type of order could intervene (this effect would have to be non-perturbative in t) before ferromagnetism develops, but the arguments given above rule out this possibility. Thus, we conclude that in the limit of small t , the singularity of the spin-triplet interaction channel leads to ferromagnetism. Our results substantiate the conclusions of previous authors^{23,24,25}.

The basic philosophy is simple: since the RG equations tell us that the interaction in the triplet channel grows in importance as the energy scale is lowered, we try to satisfy this term in the action first by allowing the magnetization, X , to develop a non-zero expectation value. In order to do this, we compute an effective action $S_{\text{eff}}[X]$ for the magnetization X taking Finkelstein's 'non-linear σ -model' for interacting electrons in a disordered system as the point of departure. Since our goal is to calculate an effective action $S_{\text{eff}}[X]$ to lowest order in the resistance t , Finkelstein's σ -model is merely a convenient way of organizing the pertinent diagrams. We believe that it is intuitively appealing to follow Finkelstein and work directly with a theory which retains only diffusion modes and drops the superfluous high-energy electronic degrees of freedom, but this is a matter of convenience, and it is an option which we can take regardless of whether new terms are generated at the two-loop or higher level (i.e. regardless of whether the non-linear terms in the theory are really related by symmetry, as they would be in a σ -model). Our derivation emphasizes the similarity between, on the one hand, the RG equation for the triplet-channel interaction in a dirty metal and the consequent development of ferromagnetism with, on the other hand, the RG equation for the Cooper-channel interaction in a clean Fermi liquid and the consequent development of

superconductivity. We believe that this illuminates the inevitability of our conclusion.

Once ferromagnetic order has developed, the system becomes essentially one of spinless electrons whose diffusion is decoupled from the spin waves of the system. In this regime, the suppression of the resistivity which accompanies the growth of the triplet interaction parameter is gone. Thus, our results indicate that a metallic state, if it exists in two dimensions, does not owe its existence to the growth of the triplet interaction parameter.

In two dimensions, our results should be interpreted as follows. As we lower the temperature, the system develops local ferromagnetic order as we cross the mean-field transition temperature T_c^{MF} . The spin density is controlled by classical thermal fluctuations which prevent order and lead to a correlation length $\xi \sim e^{(\text{const.})/T}$ at low-temperatures. In the regime $0 < T < T_c^{MF}$, we have a system in which charge diffuses while the spins are gapped. The localization physics of this system is essentially the same as that of spinless electrons, so the system leaves the diffusive regime and eventually becomes localized. Thus, our small t analysis cannot be used all the way to zero temperature. The trend is towards a ferromagnetic insulator, but the onset of insulating behavior can thwart the development of ferromagnetism. Recent experiments²⁶ performed near the metal-insulator transition indicate an enhancement of the spin susceptibility which is cut off by the onset of insulating behavior. The metal-insulator transition is outside of the regime of validity of our calculation, but this is, nevertheless, qualitatively consistent with our picture.

In $2 + \epsilon$ dimensions, when the interaction strength is larger than a threshold value of $O(\epsilon)$, there is a ferromagnetic transition at a finite temperature T_c . If the resistance is smaller than a critical value $t_c = O(\epsilon)$, then the resistance will remain finite down to zero temperature, where the system is in a metallic ferromagnetic state.

II. THE MODEL

We begin with a system of interacting electrons moving in a quenched random potential $V(x)$ in $2D$. The imaginary-time action is

$$S = S_0 + S_{\text{disorder}} + S_{\text{int}} \quad (1)$$

where

$$S_0 + S_{\text{disorder}} = T \sum_n \int d^2x \, \psi_{n,\alpha}^\dagger(x) \left(i\epsilon_n + \frac{1}{2m} \nabla^2 + \mu + V(x) \right) \psi_{n,\alpha}(x) \quad (2)$$

and

$$S_{\text{int}} = T \sum_{n,m,l} \int \left[\Gamma_1 \psi_{n,\alpha}^\dagger(\mathbf{k} + \mathbf{q}) \psi_{m,\alpha}(\mathbf{k}) \psi_{l,\beta}^\dagger(\mathbf{p} - \mathbf{q}) \psi_{l+m-n,\beta}(\mathbf{p}) + \Gamma_2 \psi_{n,\alpha}^\dagger(\mathbf{k} + \mathbf{q}) \psi_{m,\alpha}(\mathbf{p}) \psi_{l,\beta}^\dagger(\mathbf{p} - \mathbf{q}) \psi_{l+m-n,\beta}(\mathbf{k}) \right] \quad (3)$$

For later convenience, we have written the second term in momentum space and have split the interaction into nearly forward scattering, in which electrons with momenta and spins $(\mathbf{k}, \alpha)$, $(\mathbf{p}, \beta)$ are scattered to $(\mathbf{k}', \alpha)$,

$(\mathbf{p}', \beta)$, where $\mathbf{k}' \approx \mathbf{k}$, $\mathbf{p}' \approx \mathbf{p}$ and exchange scattering, in which $\mathbf{k}' \approx \mathbf{p}$, $\mathbf{p}' \approx \mathbf{k}$. If the microscopic electron-electron interaction is $V(\mathbf{q})$, then the associated bare coupling constants are $\Gamma_1 = V(0)$, $\Gamma_2 = \int (d\theta/2\pi) V(2k_F \cos \theta/2)$. As Finkelstein noted, these are subject to finite Fermi liquid renormalization in the ballistic regime, even before they begin to flow in the diffusive regime. We assume that the system is in a weak magnetic field, so that we can ignore Cooper scattering, for which $\mathbf{k}_1 \approx -\mathbf{k}_2$, $\mathbf{k}_3 \approx -\mathbf{k}_4$.

In order to perform the average over the quenched random potential, we replicate the theory and then average the replicated functional integral over the probability distribution $\overline{V(\mathbf{x})V(\mathbf{x}')} = \frac{1}{2\pi N_F \tau} \delta^{(d)}(\mathbf{x} - \mathbf{x}')$. Now, we have

$$S^{\text{rep}} = T \sum_{n;a} \int d^2x \left[\psi_{n,\alpha}^\dagger(x) \left(i\epsilon_n + \frac{1}{2m} \nabla^2 + \mu \right) \psi_{n,\alpha}(x) + \frac{1}{4\pi N_F \tau} (\psi_{n,\alpha}^\dagger \psi_{n,\alpha})^2 \right] \\ T \sum_{n,m,l;a} \int \left[\Gamma_1 \psi_{n,\alpha}^{a\dagger}(\mathbf{k} + \mathbf{q}) \psi_{m,\alpha}^a(\mathbf{k}) \psi_{l,\beta}^{a\dagger}(\mathbf{p} - \mathbf{q}) \psi_{l+m-n,\beta}^a(\mathbf{p}) + \Gamma_2 \psi_{n,\alpha}^{a\dagger}(\mathbf{k} + \mathbf{q}) \psi_{m,\alpha}^a(\mathbf{p}) \psi_{l,\beta}^{a\dagger}(\mathbf{p} - \mathbf{q}) \psi_{l+m-n,\beta}^a(\mathbf{k}) \right] \quad (4)$$

where $a = 1, 2, \dots, N$ is a replica index; eventually, we must take the replica limit, $N \rightarrow 0$. N_F is the single-particle density-of-states at the Fermi energy. The resulting effective action now contains four quartic terms: one disorder term and two interaction terms. Note that the disorder term is non-local in time since it only depends on two Matsubara frequencies, and it couples different replicas. The interaction terms depend on three Matsubara frequencies, so they are local in time, and they do not couple different replicas.

III. FINKELSTEIN'S EFFECTIVE FIELD THEORY

We decouple these quartic terms with three Hubbard-Stratonovich fields, $Q_{nm,\alpha\beta}^{ab}$, $Y_{nm}^a = Y^a(n-m)$,

$X_{nm,\alpha\beta}^a = X_{\alpha\beta}^a(n-m)$. It will be useful to think of X, Y both as matrices with indices n, m and also as functions of a single frequency ω_{n-m} ; they have only a single replica index, however, because interactions do not mix different replicas. Note that $X_{\alpha\beta} \bar{\sigma}_{\beta\alpha}$ and Y are essentially the spin and charge density. Integrating out the fermions, we can write the partition function as

$$Z = \int \mathcal{D}Q \mathcal{D}X \mathcal{D}Y e^{-S_{\text{eff}}[Q, X, Y]} \quad (5)$$

where $S_{\text{eff}}[Q, X, Y]$ is given by

$$S_{\text{eff}}[Q, X, Y] = \sum \int \left[-\text{tr} \ln \left(i\epsilon_n + \frac{1}{2m} \nabla^2 + \mu + \frac{i}{2\tau} Q + i\Gamma_1 Y + \Gamma_2 X \right) + \frac{\pi N_F}{4\tau} \text{tr} (Q^2) + \frac{1}{2} \Gamma_1 \text{tr} (Y^2) + \frac{1}{2} \Gamma_2 \text{tr} (X^2) \right] \quad (6)$$

In this expression, we use 'tr' to mean the trace over all of the indices which are not explicitly written. Matsubara indices are treated as ordinary matrix indices except that their summations come with factors of T .

The saddle-point equations are

$$\pi N_F \hat{Q} = i \int \frac{d^2p}{(2\pi)^2} \frac{1}{i\epsilon_n + \frac{1}{2m} \nabla^2 + \mu + \frac{i}{2\tau} \hat{Q} + i\Gamma_1 Y + \Gamma_2 X} \quad (7)$$

$$X(m) = T \sum_n \int \frac{d^2 p}{(2\pi)^2} \frac{1}{i\epsilon_n \delta_{n+m,n} + \frac{1}{2m} \nabla^2 + \mu + \frac{i}{2\tau} \hat{Q} + i\Gamma_1 \hat{Y} + \Gamma_2 \hat{X}} \quad (8)$$

$$Y(m) = iT \sum_n \int \frac{d^2 p}{(2\pi)^2} \frac{1}{i\epsilon_n \delta_{n+m,n} + \frac{1}{2m} \nabla^2 + \mu + \frac{i}{2\tau} \hat{Q}_{n+m,m} + i\Gamma_1 \hat{Y} + \Gamma_2 \hat{X}} \quad (9)$$

The momentum integrals on the right-hand-sides of (8) and (9) are simply $i\pi N_F Q_{n+m,n}$, according to (7). Hence, they have the solution

$$\begin{aligned} Q_{mn,\alpha\beta}^{ab} &= \text{sgn}(\epsilon_n) \delta_{mn} \delta_{\alpha\beta} \delta^{ab} \\ X_{\alpha\beta}^a(m) &= 0 \\ Y^a(m) &= 0 \end{aligned} \quad (10)$$

Note that the right-hand sides of the second and third equations of (8) and (9) vanish for $m \neq 0$ because the saddle-point Q is diagonal in Matsubara frequency while they vanish for $m = 0$ because of cancellation between positive and negative frequencies. As we will see later, for Γ_2 sufficiently large, $X = 0$ becomes unstable and there is an $X \neq 0$ solution of (8). This is the Stoner instability. Upon including low-energy fluctuations of Q – diffusion modes – an $X \neq 0$ solution arises even for Γ_2 small, as we will see shortly.

The low-energy fluctuations about this saddle-point are transverse fluctuations of Q , which can be parametrized by $V_{nm,\alpha\beta}^{ab}$:

$$Q = \begin{pmatrix} (1 - VV^\dagger)^{1/2} & V \\ V^\dagger & - (1 - V^\dagger V)^{1/2} \end{pmatrix} \quad (11)$$

The diagonal blocks of Q_{nm} correspond to Matsubara indices $n > 0, m > 0$ and $n < 0, m < 0$ while the upper right block corresponds to $n < 0, m > 0$; the lower left block, to $n > 0, m < 0$. These fluctuations correspond to diffusion of charge and spin. The longitudinal fluctuations of Q as well as fluctuations of X, Y are gapped at the classical level (i.e. tree-level).

In order to derive an effective field theory for these diffusion modes, we shift $Q \rightarrow Q - 2\tau\Gamma_1 Y + 2\tau i\Gamma_2 X$ in (6)

$$\begin{aligned} S_{\text{eff}}[Q, X, Y] = \sum \int & \left[-\text{tr} \ln \left(i\epsilon_n + \frac{1}{2m} \nabla^2 + \mu + \frac{i}{2\tau} Q \right) + \frac{\pi N_F}{4\tau} \text{tr} (Q^2) - \pi N_F \Gamma_1 \text{tr} (QY) + i\pi N_F \Gamma_2 \text{tr} (QX) \right. \\ & \left. + \frac{1}{2} \left(1 + 2\pi \left(\tau T \sum_m \right) N_F \Gamma_1 \right) \Gamma_1 \text{tr} (Y^2) + \frac{1}{2} \left(1 - 2\pi \left(\tau T \sum_m \right) N_F \Gamma_2 \right) \Gamma_2 \text{tr} (X^2) \right] \end{aligned} \quad (12)$$

Formally, the factors of $\tau(T\sum_m)$ in the final two terms are infinite as a result of the unrestricted Matsubara sum. However, upon integrating out the massive longitudinal modes of Q , we see that they are the first terms in the series $\tau T \sum_m - 2\pi(\tau T \sum_m)^2 + \dots = \tau T \sum_m / (1 + 2\pi\tau T \sum_m)$, which is simply $1/2\pi$. Thus, we replace the factors of $\tau(T\sum_m)$ by $1/2\pi$ in (12). Alternatively, we could expand the $\text{tr} \ln(\dots)$ about the diffusive saddle-point and keep only terms up to second order in X and Y . We would then obtain the same expressions, namely (12) with the factors of $\tau(T\sum_m)$ replaced by $1/2\pi$. The coefficient of X^2 is now $1 - N_F \Gamma_2$, which becomes negative for $\Gamma_2 > 1/N_F$: the Stoner instability. Finkelstein actually expanded the $\text{tr} \ln(\dots)$ only to linear order in X and Y . For small Γ_1, Γ_2 , there is no difference, but the Stoner instability is missed.

We now expand the $\text{tr} \ln(\dots)$ about the diffusive saddle-point. Thus, we require that Q is a constrained field of the form given in (11).

$$\begin{aligned} S_{\text{eff}}[Q] = \pi N_F \int d^d x & \left\{ D \text{tr} (\nabla Q)^2 - 4iZ \text{tr} (\hat{\epsilon} Q) \right. \\ & - \Gamma_1 \text{tr} (QY) + i\Gamma_2 \text{tr} (QX) \\ & + \frac{1}{2} (1 + N_F \Gamma_1) \frac{\Gamma_1}{\pi N_F} \text{tr} (Y^2) \\ & \left. + \frac{1}{2} (1 - N_F \Gamma_2) \frac{\Gamma_2}{\pi N_F} \text{tr} (X^2) \right\} \end{aligned} \quad (13)$$

The diffusion constant D is given by $D = v_F^2 \tau / 2$.

Following Finkelstein, we integrate out X, Y and obtain an effective action for the diffusion modes.

$$S_{\text{eff}}[Q] = \pi N_F \int d^d x \left\{ D \text{tr}(\nabla Q)^2 - 4i \text{tr}(\hat{\epsilon} Q) \right. \\ \left. - \pi N_F \tilde{\Gamma}_1 Q_{n_1 n_2, \alpha \alpha}^{aa} Q_{n_3 n_4, \beta \beta}^{aa} \delta_{n_1 - n_2 + n_3 - n_4} \right. \\ \left. + \pi N_F \tilde{\Gamma}_2 Q_{n_1 n_2 \alpha \beta}^{aa} Q_{n_3 n_4 \beta \alpha}^{aa} \delta_{n_1 + n_2 - n_3 - n_4} \right\} \quad (14)$$

Note that ordinary matrix multiplication rules reflect the non-locality in time of the first term (the ‘disorder term’) in this action. The $\tilde{\Gamma}_{1,2}$ interaction terms do not involve matrix multiplication and are, consequently, local in time. In this expression, $\tilde{\Gamma}_{1,2} = \Gamma_{1,2}/(1 \pm N_F \Gamma_{1,2})$. These corrections to $\Gamma_{1,2}$ follow from our retention of the X^2 and Y^2 terms which Finkelstein drops. For $\Gamma_{1,2}$ small, $\tilde{\Gamma}_{1,2} = \Gamma_{1,2}$, and Finkelstein’s effective action is recovered.

Initially, the coefficient of the $\text{tr}(\hat{\epsilon} Q)$ term is 1, as in (13), but quantum corrections cause it to flow, so we have followed Finkelstein in introducing the coupling Z .

IV. RENORMALIZATION GROUP FLOWS

The one-loop RG equations associated with this action are

$$\begin{aligned} \frac{dt}{d\ell} &= t^2 \left(4 - 3 \frac{Z_2}{\Gamma_2} \ln \left(\frac{Z_2}{Z} \right) + \frac{Z_1}{2\Gamma_1 - \Gamma_2} \ln \left(\frac{Z_1}{Z} \right) \right) \\ \frac{dZ}{d\ell} &= t(-\Gamma_1 + 2\Gamma_2) \\ \frac{d\Gamma_1}{d\ell} &= t \left(\Gamma_2 + \frac{\Gamma_2^2}{Z} \right) \\ \frac{d\Gamma_2}{d\ell} &= t \left(\Gamma_1 + \frac{2\Gamma_2^2}{Z} \right) \end{aligned} \quad (15)$$

where t is the dimensionless resistance, $t = 1/(2\pi D)$; $Z_1 = Z - 2\Gamma_1 + \Gamma_2$; $Z_2 = Z + \Gamma_2$; and we have set $\pi N_F = 1$ in order to avoid clutter.

For sufficiently large interaction strength, the correct starting point is presumably a Wigner crystal, which corresponds to a radically different saddle-point of the effective action (6). Thus, even though these RG equations were computed to all orders in $\Gamma_{1,2}$ for t small, they should be understood as holding only for the vicinity of the diffusive saddle-point.

The RG equations for Z , Γ_1 , Γ_2 can be combined to give:

$$\frac{d}{d\ell} (Z - 2\Gamma_1 + \Gamma_2) = 0 \quad (16)$$

Hence, $Z - 2\Gamma_1 + \Gamma_2$ is a constant. For long-ranged Coulomb interactions, the Ward identity implies that this constant is zero: $Z - 2\Gamma_1 + \Gamma_2 = 0$. More generally, we can introduce $\gamma_1 = \Gamma_1/Z$, $\gamma_2 = \Gamma_2/Z$, so that $\gamma_1 = \frac{1}{2}(1 + \gamma_2) - \text{const.}/Z$. Since Z grows under the RG

flow (15), the final term can be dropped and we have $\gamma_1 = (1 + \gamma_2)/2$. In the case of spinless electrons, Z does not grow under the corresponding RG flow, so we cannot make this simplification; we will later discuss the similar case of spin-polarized electrons.

Making the above substitution, we find that the RG equations for γ_2 and Z are:

$$\begin{aligned} \frac{d\gamma_2}{d\ell} &= t(1 + \gamma_2)^2 \\ \frac{dZ}{d\ell} &= Z t \left(-\frac{1}{2} + \frac{3}{2} \gamma_2 \right) \end{aligned} \quad (17)$$

Thus, γ_2 and Z grow as the scale is increased. Eventually, this growth causes these equations to break down. If we ignore the flow of t , then the breakdown scale is $E \sim (1/\tau) e^{-\frac{1}{t(1+\gamma_2)}}$; in fact, the scale is somewhat lower because t flows.

As noted in ref. 7, the breakdown of these equations is reminiscent of the situation in a clean Fermi liquid, where the Cooper channel interaction flows according to $dV/d\ell = -V^2$. The breakdown of the latter at a scale $\Delta \sim \Lambda e^{-1/V}$ signals the onset of superconductivity or, at least, pairing. Unlike in the case of a clean Fermi liquid with a four-fermion Cooper channel interaction, Z diverges as well in our problem. However, the divergence of Z is a by-product of the divergence of γ_2 ; In the Cooper pairing problem, if the interaction which drives pairing could also contribute to mass renormalization, we would have the same situation as in the Finkelstein problem. Indeed, this is what happens when pairing is driven by a gauge field. The development of pairing not only cuts off the flow of the interaction, but also the flow of the effective mass²⁷. Similarly, we expect that the physics which resolves the divergence of γ_2 will also resolve the divergence of Z .

V. EFFECTIVE ACTION FOR MAGNETIC DEGREES OF FREEDOM: ONE-LOOP

To see how the breakdown of (15) leads to ferromagnetism, let us go back to the action (13). Instead of integrating out X and Y at the paramagnetic diffusive saddle point, let’s integrate out V and Y in order to obtain an effective action $S_{\text{eff}}[X]$ for the magnetization X .

The spin polarization X acts as a Zeeman field. A constant field aligned along the z direction, $X_{\alpha\beta}^a = X \sigma_{\alpha\beta}^z$ (taken to be replica-diagonal), gives a gap to $V_{\uparrow\downarrow}$ and $V_{\downarrow\uparrow}$ but leaves $V_{\uparrow\uparrow}$ and $V_{\downarrow\downarrow}$ unaffected. To see this, note that a constant field couples only to the diagonal Matsubara indices Q_{nn} . Expanding $(1 - VV^\dagger)^{1/2}$ and $-(1 - V^\dagger V)^{1/2}$ and retaining only the leading terms, which are quadratic, we find that

$$i\Gamma_2 \text{tr}(XQ) = i\Gamma_2 X \text{tr} \left[\left(V_{\uparrow\uparrow} V_{\uparrow\downarrow}^\dagger - V_{\uparrow\downarrow} V_{\uparrow\uparrow}^\dagger \right) \right] \quad (18)$$

An imaginary gap $\pm i\Gamma_2 X$ in Matsubara formalism corresponds to a real gap in real-time. $V_{\uparrow\downarrow}$ and $V_{\downarrow\uparrow}$ have

propagators:

$$\left\langle V_{nm; \uparrow\downarrow, \uparrow\downarrow}(\mathbf{q}) V_{nm; \downarrow\uparrow, \downarrow\uparrow}(-\mathbf{q}) \right\rangle = \frac{1}{Z|\omega_{nm}| + Dq^2 \pm i\Gamma_2 X} \quad (19)$$

When we integrate out $V_{\uparrow\downarrow}$ and $V_{\downarrow\uparrow}$, infrared divergences will be cutoff by X , as the single-particle gap cuts off divergences in BCS theory. Since $V_{\uparrow\uparrow}$ and $V_{\downarrow\downarrow}$ are unaffected by X to this order, they simply contribute an unimportant constant to $S_{\text{eff}}[X]$.

We will integrate out V to one loop to obtain $S_{\text{eff}}[X]$.

This is not quite the same as a calculation to lowest order in t , but the basic physics is illustrated most simply in a one-loop calculation, so we will present it first. To maintain further simplicity, let us consider the case $\Gamma_1 = 0$; the general case is a straightforward but more complicated extension, to which we return later. In the next subsection, we will present a more careful calculation which includes the effects of Γ_1 and includes all of the terms which contribute to lowest order in t .

At one loop, we only need to retain terms in (13) up to quadratic order in V

$$S[V, X] = \pi N_F \sum \int \left\{ D \text{tr} (\nabla V_+^\dagger \nabla V_+) + Z|\omega_{nm}| V_{nm,+}^\dagger V_{mn,+} + \frac{1}{2} (1 - N_F \Gamma_2) \frac{\Gamma_2}{\pi N_F} \text{tr} (X^2) \right. \\ \left. + i\Gamma_2 [\text{tr} (V_{nm,+} X(n-m)) + \text{c.c.}] + i\Gamma_2 [\text{tr} (V_{ml,+} V_{ln,+}^\dagger X(n-m)) + \text{c.c.}] + (V_+ \rightarrow V_-, X \rightarrow -X) \right\} \quad (20)$$

Here, we have written $V_+ \equiv V_{\uparrow\downarrow}$, $V_+^\dagger \equiv V_{\downarrow\uparrow}^\dagger$, $V_- \equiv V_{\downarrow\uparrow}$, and $V_-^\dagger \equiv V_{\uparrow\downarrow}^\dagger$.

We now shift X in order to remove the bi-linear coupling between X and V . This is the simplest way to proceed with a one-loop computation of $S_{\text{eff}}[X]$. To go beyond one-loop, it is more convenient to shift V , as we will see later. Performing the shift $X(l) \rightarrow X(l) - iT \sum_{m,A} V_{m+l,m;A}$ for $l > 0$ and $X(l) \rightarrow X(l) - iT \sum_{n,A} V_{n+l,n;A}^\dagger$ for $l < 0$, where $A = \pm$, we have:

$$S[V, X] = \pi N_F \sum \int \left\{ D \text{tr} (\nabla V^\dagger \nabla V) + Z|\omega_{nm}| V_{nm;A}^\dagger V_{mn;A} - \Gamma_2 V_{nm;A}^\dagger V_{m'n';A} \delta_{n-m,n'-m'} \right. \\ \left. + i\Gamma_2 [\text{tr} (V_{ml;A} V_{ln;B}^\dagger \sigma_{AB}^z X(n-m)) + \text{c.c.}] + \frac{1}{2} (1 - N_F \Gamma_2) \frac{\Gamma_2}{\pi N_F} \text{tr} (X^2) \right\} \quad (21)$$

A V^3 term is also generated as a result of this shift, but it doesn't contribute to $S_{\text{eff}}[X]$ at one loop, so we drop it. The V propagator which follows from (21) is now

$$\left\langle V_{nm; \pm; ab}(\mathbf{q}) V_{nm; \pm; ba}(-\mathbf{q}) \right\rangle = D_0^{X\pm}(\omega_{nm}, q) + \Gamma_2 D_0^{X\pm}(\omega_{nm}, q) D_2^{X\pm}(\omega_{nm}, q) \delta_{ab} \quad (22)$$

where

$$D_0^{X\pm}(\omega_{nm}, q) = \frac{1}{Z|\omega_{nm}| + Dq^2 \pm i\Gamma_2 X} \\ D_2^{X\pm}(\omega_{nm}, q) = \frac{1}{Z_2|\omega_{nm}| + Dq^2 \pm i\Gamma_2 X} \quad (23)$$

and Z_2 is as defined after (15).

Integrating out V , we obtain all one-loop diagrams of (21), with X treated as a classical background field or, simply, a $\text{tr} \ln(\dots)$ expression containing the V propagators in the X background field. Later, we will return to the possibility of other saddle-points, but for now, let us assume, in the name of simplicity, that X is static, translationally-invariant, and replica-symmetric.




FIG. 1: The one-loop diagram with X -treated as a classical background field can be viewed as a sum over one-loop diagrams with X insertions (the thick lines on the right).

For such X , the effective action takes the form:

$$S_{\text{eff}}[X] = \sum \int \frac{1}{2} (1 - N_F \Gamma_2) \Gamma_2 \text{tr} (X^2) \\ + \text{tr} \ln \left([Z|\omega_{nm}| + Dq^2 \pm i\Gamma_2 X] \delta_{nn'} \delta_{mm'} \delta_{aa'} \delta_{bb'} \right. \\ \left. - \Gamma_2 \delta_{n-m-n'+m'} \delta_{aa'} \delta_{bb'} \delta_{ab} \right) \quad (24)$$

This may be rewritten in terms of the propagators $D_{0,2}^{X\pm}$

as

$$S_{\text{eff}}[X] = \sum \int \frac{1}{2} (1 - N_F \Gamma_2) \Gamma_2 \text{tr} (X^2) + \text{tr} \ln \left[(D_0^{X+} - \Gamma_2 D_0^{X+} D_2^{X+} \delta_{ab}) (D_0^{X-} - \Gamma_2 D_0^{X-} D_2^{X-} \delta_{ab}) \right] \quad (25)$$

When we take the product of the two factors on the right-hand side of (25), the $D_0^{X+} D_0^{X-}$ term has a free summation over the replica index b which vanishes in the replica limit $N \rightarrow 0$ for replica-symmetric solutions $X_a = X$. Dropping this term and using the relation $D_0^{X\pm} - \Gamma_2 |\omega_{nm}| D_0^{X\pm} D_2^{X\pm} = D_2^{X\pm}$, we have

$$S_{\text{eff}}[X] = \sum \int \left\{ \frac{1}{2} (1 - N_F \Gamma_2) \Gamma_2 \text{tr} (X^2) - \text{tr} \ln (D_2^{X+} D_2^{X-} - D_0^{X+} D_0^{X-}) \right\} \quad (26)$$

Evaluating the integral, we find

$$S_{\text{eff}}[X] = \beta L^2 \left[\frac{1}{2} (1 - N_F \Gamma_2) \Gamma_2 X^2 - t \pi N_F \Gamma_2^2 \left(\frac{1}{Z} - \frac{1}{Z_2} \right) X^2 \left(\ln \left(\frac{\Lambda}{\Gamma_2 X} \right) - 1 \right) \right] \quad (27)$$

where L^2 is the area of the system.

As a result of the logarithm on the second line, the minimum of this effective action is at non-zero X for arbitrarily small Γ_2 . We will discuss this non-trivial solution and the saddle-point equation (or gap equation) which it satisfies in more detail in the next section. At this stage, the important lesson is that, at the one-loop level, the weakly-interacting diffusive electron gas is always unstable to the development of non-zero X , i.e. ferromagnetism.

VI. FERROMAGNETIC SADDLE-POINT AT ONE-LOOP

Let us examine more closely the saddle-point equation for X in the simplest setting, that of section V. If we vary equation (25) with respect to X (before performing the momentum integral and Matsubara sum), then we obtain the saddle-point equation:

$$\begin{aligned} 1 - N_F \Gamma_2 &= \pi N_F \Gamma_2 \sum \int \left[D_2^{X+} D_2^{X-} - D_0^{X+} D_0^{X-} \right] \\ &= \pi N_F \Gamma_2 \sum \int \left[\frac{1}{[Z |\omega_{nm}| + D q^2]^2 + \Gamma_2^2 X^2} - \frac{1}{[Z |\omega_{nm}| + D q^2]^2 + \Gamma_2^2 X^2} \right] \\ &= t \pi N_F \Gamma_2 \left(\frac{1}{Z} - \frac{1}{Z_2} \right) \ln \left(\frac{\Lambda}{\Gamma_2 X} \right) \end{aligned} \quad (28)$$

The integrals on the right-hand-side of the second line are, for $X = 0$, precisely the integrals which determine the contributions of diagrams (d) and (e) to the RG equation for Γ_2 . We will further discuss this point below.

The right-hand-side of the saddle-point equation is logarithmically divergent in the $X \rightarrow 0$ limit. Hence, there is always a solution

$$\Gamma_2 X = \frac{1}{\tau} e^{-(1 - N_F \Gamma_2)(1 + \gamma_2)/(t \gamma_2^2)} \quad (29)$$

We have taken $\Lambda = 1/\tau$ as the ultraviolet cutoff and $\gamma_2 = \pi N_F \Gamma_2 / Z$. This is the non-trivial minimum of the action to which we referred at the end of section V. The structure of this saddle-point equation is clearly reminis-

cent of the BCS gap equation. The integrals on the right-hand-side are logarithmically divergent – as expected for a marginally relevant interaction. A symmetry-breaking order parameter must develop in order to cut them off.

We can submit our saddle-point solution to an important check by verifying its consistency with the RG equations. The gap to spin-flip excitations, $\Gamma_2 X$ is an observable quantity, and should be independent of the ultraviolet cutoff, Λ . Even though the cutoff is taken to be a physical scale $1/\tau$, we are free to integrate out short-wavelength fluctuations in order to lower the cutoff while simultaneously renormalizing the couplings Γ_1 , Γ_2 , t , and Z . This must occur in such a way that the saddle-point solution, $\Gamma_2 X$, remains unchanged. For this

reason, we regard the one-loop calculation of section V as incomplete. It is consistent with an RG equation which only includes diagrams (d) and (e) of figure 2 for the flow of Γ_2 . Diagrams (a), (b), and (c) are not incorporated, and neither is the flow of Γ_1 , t , Z .




FIG. 2: The five basic diagrams which contribute to the renormalization of Γ_2 . Solid lines are V propagators. Dotted lines are bare X propagators.

VII. BEYOND ONE-LOOP

Though the preceding one-loop calculation captures the basic physics of the problem, it does not capture all of the order t physics. As we have just noted, the effective action (24) only includes terms corresponding to two of the five diagrams in figure 2 – namely, (d) and (e) – which contribute to (15).

The apparent contradiction stems from the fact that the physics of a one-loop RG equation might only appear at higher-loops in a saddle-point calculation. This is surprising at first glance, but can be understood at a technical level by noting that a contribution to the effective action of the form $t^2 \ln^2(\Lambda/X)$ is equally important at the saddle-point as a contribution of the form $t \ln(\Lambda/X)$ since the extra logarithm compensates for the extra factor of t . A simple example of a model in which this occurs is given in the appendix.

The effects of Γ_1 on the flow of Γ_2 , t , Z enter in a two-loop calculation of the effective action. Consider the diagrams of figure 3. These are two-loop diagrams which, when combined with the one-loop diagram of the previous section, have the effect of shifting in the effective action of the previous section the values of Γ_2 , t , and Z from their bare values. These diagrams can receive contributions from non-spin-flip diffusion propagators which are not cut off by the magnetization, X . They are charge and S_z diffusion modes. Since we expect them to be a part of the low-energy theory, we can choose not to integrate them out or to integrate out modes above a cutoff of our choosing. The natural choice is to integrate out these modes above $\Gamma_2 X$ so that the resulting effective field theory for X , $V_{\uparrow\uparrow}$, $V_{\downarrow\downarrow}$ contains only excitations with energies less than $\Gamma_2 X$. (We could alternatively imagine performing the calculation at finite-temperature so

that T is effectively the cutoff for both spin-flip and non-spin-flip propagators. Such a calculation could be used to obtain the transition temperature T_c for the putative transition.) This effects the following replacements:

$$\begin{aligned}\Gamma_2 &\rightarrow \Gamma_2 + t\Gamma_1 \ln\left(\frac{\Lambda}{\Gamma_2 X}\right) \\ t &\rightarrow t + t^2\Gamma_1 f_2(z, z_1, z_2) \ln\left(\frac{\Lambda}{\Gamma_2 X}\right) \\ Z &\rightarrow Z - t\Gamma_1 \ln\left(\frac{\Lambda}{\Gamma_2 X}\right)\end{aligned}\quad (30)$$

where

$$\begin{aligned}f_2(z, z_1, z_2) &= \frac{2z_1}{z_1 - z_2} f_1(z, z_1) - \frac{2z_2}{z_1 - z_2} f_1(z, z_2) \\ f_1(z, z_2) &= \frac{1}{z_2 - z} \ln\left(\frac{z_2}{z}\right)\end{aligned}\quad (31)$$




FIG. 3: Two-loop contributions to $S_{\text{eff}}[X]$ which renormalize Γ_2 , t , and z . The first diagram renormalizes Γ_2 while the second renormalizes t and z . The quadratic vertices are Γ_1 interaction vertices.

Diagrams (a)-(c) of figure 2 cannot contribute directly to $S_{\text{eff}}[X]$ for static X because the external V lines cannot carry zero frequency. The Matsubara indices m , n of V_{mn} necessarily satisfy $n > 0$, $m < 0$, so their difference cannot vanish. For the same reason, these diagrams cannot be linked with (d) and (e), in the way demonstrated in the toy model in the appendix. It is possible that contributions related to diagrams (a)-(c) enter $S_{\text{eff}}[X]$ through even higher-loop contributions or, perhaps, when fluctuations in X are included. However, this touches upon the unresolved issue of the renormalizability of the theory at higher-loops. In the absence of a clearer understanding of such higher-loop contributions to $S_{\text{eff}}[X]$, we conclude that the one-loop effective action (27) with the replacement (30) is the most systematic result which we can obtain at present. We expect that going beyond one-loop in V or including the effects of fluctuations of X will quantitatively but not qualitatively modify our result.

In section V, we removed the bi-linear coupling between X and V by shifting X . It may seem more natural to remove the bi-linear coupling between X and V by performing the shift $V_{nm} \rightarrow V_{nm} - D_0^{-1}(q, \omega_{nm})X(n-m)$. If we then integrate out V , we will not find a non-trivial replica-symmetric saddle-point (though there may

be replica-symmetry-breaking saddle-points), and the resulting action will not contain any of the physics of the one-loop RG equations. In order to recover this physics, we must now go to one-loop in $S_{\text{eff}}[X]$. Going to one-loop in $S_{\text{eff}}[X]$, we recover the saddle-point of our initial action of section V. The relevant diagrams are diagrams (d) and (e) of figure 2, but the solid lines are shrunk to a point when these diagrams are viewed as 1-loop diagrams in $S_{\text{eff}}[X]$. Diagrams corresponding to (a), (b), and (c) have dotted lines attached to the solid external legs (via the bi-linear coupling between X and V); however, these vanish in the static limit for the same reason as above. Thus, at this level, there is no advantage in proceeding this way, and there is a definite disadvantage in that one extra step is necessary. However, it is possible that a

higher-loop calculation is simpler in this approach.

The one-loop effective action which we computed in section V is actually exact in a generalization of the model. The most naive large- N version of the model is not particularly simple. If we simply introduce N flavors of electrons ψ_α , $\alpha = 1, \dots, N$, then the Q -matrix has two flavor indices, Q_{AB} , and the large- N limit allows us to drop all non-planar diagrams. However, we do not know how to sum all of the planar diagrams. Instead, we consider the following generalization: we suppose that there are N flavors of diffuson fields V_α , $\alpha = 1, \dots, N$. This generalization does not have a simple interpretation in terms of the original electron variables, but it is technically simple. The action (21) is generalized to:

$$S[V, X] = \pi N_F \sum \int \left\{ D \text{tr} (\nabla V^\dagger \nabla V) + Z |\omega_{nm}| V_{nm; A\alpha}^\dagger V_{mn; A\alpha} - \Gamma_2 V_{nm; A\alpha}^\dagger V_{m'n'; A\alpha} \delta_{n-m, n'-m'} \right. \\ \left. + i\Gamma_2 \left[\text{tr} \left(V_{ml; A\alpha} V_{ln; B\alpha}^\dagger \sigma_{AB}^z X(n-m) \right) + \text{c.c.} \right] + \frac{N}{2} (1 - N_F \Gamma_2) \frac{\Gamma_2}{\pi N_F} \text{tr} (X^2) - \frac{1}{N} \Gamma_2 \sum_\beta \left[\text{tr} (V_\alpha V_\alpha^\dagger V_\beta) + \text{c.c.} \right] \right\} \quad (32)$$

Note that there is still only a single X field. The explicit factor of N in front of the penultimate term and the factor of $1/N$ in front of the final term ensures that every term in the action is $O(N)$. For this action, the diagram which we calculated in section V is the sole $O(N)$ diagram. All others are $O(1)$ or smaller. Furthermore, our effective action is multiplied by a coefficient N . Thus, in the $N \rightarrow \infty$ limit, our one-loop saddle-point calculation is exact.

At first glance, it may seem almost contradictory to take an $N \rightarrow \infty$ limit while simultaneously taking a replica limit $n \rightarrow 0$. However, the flavor index structure in the action (32) is very different from the replica index structure. The latter is chosen to ensure that closed fermion loops vanish while the latter selects a preferred set of the remaining diagrams. Hence, the two limits are consistent.

VIII. PHYSICS OF THE FERROMAGNETIC DIFFUSIVE FERMION LIQUID

Having concluded that the divergence of the triplet interaction amplitude leads to ferromagnetism, the obvious next question is “so then what?” To answer this, we must examine the diffusion of electrons in a ferromagnetic metal, and the interaction between diffusion modes and spin waves.

In the ferromagnetic state, we can expand X about its perfectly-ordered ground state, which we take to be

along the $\hat{\mathbf{z}}$ -direction

$$\mathbf{X} = \left(m_x, m_y, \sqrt{1 - m_x^2 - m_y^2} \right) \quad (33)$$

We can also perform a gradient expansion of $S_{\text{eff}}[\mathbf{X}]$. The result is essentially dictated by symmetry:

$$S_{\text{eff}}[\mathbf{X}] = \int \left(\epsilon_{ij} m_i \partial_\tau m_j + K (\nabla m_i)^2 + O((\nabla m)^4) \right) \quad (34)$$

where $i = 1, 2$ are the $\hat{\mathbf{x}}$ and $\hat{\mathbf{y}}$ components of $\mathbf{X}$. The constant K is the spin stiffness of the ferromagnet.

We now reinstate the coupling of the order parameter to the diffusion modes of the system, as in (13) or (20).

$$S[V, m] = \pi N_F \sum \int \left\{ D \text{tr} (\nabla V^\dagger \nabla V) + Z |\omega_{nm}| V_{nm}^\dagger V_{mn} \right. \\ \left. + i\Gamma_2 X_0 \left[\text{tr} (\sigma_z V_{nm} V_{mn}^\dagger) + \text{c.c.} \right] \right. \\ \left. + i\Gamma_2 \left[m_i (\omega_{n-m}) \text{tr} (\sigma_i V_{nm}) + \text{c.c.} \right] \right\} \\ + \sum_n \int \left(\epsilon_{ij} i\omega_n m_i m_j + K (\nabla m_i)^2 \right) \quad (35)$$

where σ_i are Pauli matrices and we have dropped higher-order terms in m_i . A purist may object that the V 's should not appear in the action for m_i since the dynamics for the order parameter was obtained by integrating out the V 's. If we now calculate with this action, we run the risk of double-counting the degrees of freedom of the

system. In order to avoid this, we must either calculate only to tree-level in V (but allow loops in m_i) or introduce counter-terms which cancel the final line of (35). In the second approach, by computing loops in V , we will cancel the m_i counterterms.

The physics of this model is not completely manifest in the above form (35) because a uniform rotation of the ferromagnetic order parameter appears to be an excitation – since there are non-derivative terms in m_i – when, in fact, it isn't. This can be made manifest by rotating the quantization axis of the diffusing electrons so that it is aligned along the local direction of the order parameter. This is done by enacting the variable change

$$V \rightarrow e^{i\epsilon_{kl}m_k\sigma_l} V e^{-i\epsilon_{ij}m_i\sigma_j} \quad (36)$$

The action is now

$$\begin{aligned} S[V, m] = N_F \sum \int \Big\{ & D \operatorname{tr} (\nabla V^\dagger \nabla V) + Z |\omega_{nm}| V_{nm}^\dagger V_{mn} \\ & + iX_0 \Gamma_2 [\operatorname{tr} (\sigma_z V V^\dagger) + \text{c.c.}] \\ & D \operatorname{tr} (\nabla V_{nm}^\dagger \nabla (\epsilon_{ij} m_i (\omega_{n-l}) [\sigma_j, V_{ml}])) + \text{c.c.} \Big\} \\ & + \sum_n \int \left(\epsilon_{ij} i \omega_n m_i m_j + K (\nabla m_i)^2 \right) \quad (37) \end{aligned}$$

As noted earlier, there is a gap for spin-flip diffusion modes, as given by the second line of (37). Thus, we can ignore these modes at low energies and focus on charge diffusion and S_z diffusion; $S_{x,y}$ are carried by spin waves $m_{x,y}$. From the structure of the interaction term on the third line of (37), we see that $m_{x,y}$ cannot couple directly to charge or S_z diffusion alone, as dictated by rotational invariance. A charge or S_z diffusion mode, upon emitting or absorbing a spin wave, must then become a spin-flip diffusion mode. Thus, this coupling can be ignored at energies below the gap for spin-flip diffusion modes. Even in the absence of such a gap, the coupling would have dimension $d/2$ in d -dimensional space and be strongly irrelevant, as expected for interactions with Goldstone modes – which must occur through derivative couplings such as that on the third line of (37).

Thus, the low-energy physics of the ferromagnetic diffusive metal state is that of ferromagnetic spin waves decoupled from diffusion in a Zeeman field. The latter has been well-studied^{3,5}, and is similar to the spinless electron problem. Interactions suppress the conductivity, as may be seen from the RG equations:

$$\begin{aligned} \frac{dt}{d\ell} &= t^2 \left(2 - \frac{Z - \Gamma_H - \Gamma_1}{\Gamma_1 + \Gamma_H} \ln \left(\frac{Z}{Z - \Gamma_H - \Gamma_1} \right) \right. \\ &\quad \left. - \frac{Z + \Gamma_H + \Gamma_1}{\Gamma_1 - \Gamma_H} \ln \left(\frac{Z - \Gamma_H + \Gamma_1}{Z} \right) \right) \\ \frac{dZ}{d\ell} &= -t\Gamma_H, \quad \frac{d\Gamma_1}{d\ell} = 0, \quad \frac{d\Gamma_H}{d\ell} = -t\Gamma_H \quad (38) \end{aligned}$$

Γ_H is the part of the interaction between $S_z = 0$ diffusion modes which includes a contribution from both the direct

and exchange channels, $\Gamma_H (Q_{\uparrow\uparrow} Q_{\uparrow\uparrow} + Q_{\downarrow\downarrow} Q_{\downarrow\downarrow})$; its bare value is $\Gamma_H = \Gamma_1 - \Gamma_2$.

In two dimensions, these RG equations describe a flow towards insulating behavior. However, there is a metal-insulator transition in $2 + \epsilon$ dimensions through which the system can be tuned by varying the bare resistivity through a critical value of $O(\epsilon)$. We examine the situation in some more detail in the next section.

The preceding discussion is strictly true at zero temperature. At finite temperatures, thermal fluctuations will destroy two-dimensional long-range ferromagnetic order, but some signature of the zero-temperature ferromagnetic phase persists. If a finite external magnetic field is applied, the magnetization is a scaling function of temperature and magnetic field. A large- N calculation for an $SU(N)$ magnet theory gives the following expression for the scaling function for the susceptibility²⁸:

$$\chi(r, h) = \frac{M_0}{8\pi r} \ln \left[(q - e^{-h/2}) / (q - e^{h/2}) \right] \quad (39)$$

where $r = K/T$ and $h = H/T$ are the rescaled spin stiffness and magnetic field respectively, q is a root of the equation $(q - e^{-h/2})(q - e^{h/2}) = q^2 e^{-8\pi r}$, and M_0 is the zero temperature magnetization. The susceptibility is given by $M_0(e^{4\pi r} - 1)/(8\pi T r)$, which diverges exponentially as $T \rightarrow 0$.

In the limit of weak interactions and weak disorder, there is a substantial temperature regime in which there is ferromagnetic order with a long – but finite – correlation length and the resistivity is small. If the dimensionless resistivity, t , is taken to be 0.3 at temperature 5 K, and r_s , the average interspace between electrons measured in the unit of effective Bohr radius, is taken to be 0.1, then the susceptibility is large below 15 mK. However, the resistivity becomes $O(1)$ at temperatures $\ll 10^{-20}$ K. While ferromagnetism sets in at very low temperatures, insulating behavior is completely academic. In the temperature interval between these two scales, we expect the susceptibility to be of the above form (39). When r_s is larger, the separation between these scales decreases, but we should also be more circumspect about using this weak-coupling theory. Of course, if the magnetic field is removed, we pass from the unitary ensemble to the orthogonal one, and weak-localization effects should drive the system insulating much more rapidly.

IX. PHASE DIAGRAM IN $2 + \epsilon$ DIMENSIONS

In $2 + \epsilon$ dimensions, the renormalization group equations are modified by the appearance of tree-level dimensions associated with the spatial dimensionality:

$$\begin{aligned} \frac{dt}{d\ell} &= -\epsilon t + t^2 \left(4 - 3 \frac{Z_2}{\Gamma_2} \ln \left(\frac{Z_2}{Z} \right) + \frac{Z_1}{2\Gamma_1 - \Gamma_2} \ln \left(\frac{Z_1}{Z} \right) \right) \\ \frac{dZ}{d\ell} &= t(-\Gamma_1 + 2\Gamma_2) \end{aligned}$$

$$\begin{aligned}\frac{d\Gamma_1}{d\ell} &= t \left(\Gamma_2 + \frac{\Gamma_2^2}{Z} \right) \\ \frac{d\Gamma_2}{d\ell} &= t \left(\Gamma_1 + \frac{2\Gamma_2^2}{Z} \right)\end{aligned}\quad (40)$$

By varying Γ_2 , we can tune the system through a paramagnet-ferromagnet transition.

For t, γ_2 small, the flows are into the fixed line $t = 0$. This fixed line terminates at $\Gamma_2 = \Gamma_2^{\text{Stoner}}$. For $\Gamma_2 = 0$, there is a metal-insulator transition at $t = \epsilon$. In the vicinity of this transition, γ_2 is relevant. If it flows to sufficiently large values, then ferromagnetism will develop. However, if t flows to zero before this happens, then the system will remain paramagnetic. Both possibilities can occur as soon as γ_2 becomes nonzero, so the ferromagnet-paramagnet boundary which begins on the $t = 0$ axis at $\Gamma_2 = \Gamma_2^{\text{Stoner}}$ terminates at $t = \epsilon, \gamma_2 = 0$. Ferromagnetism, if it occurs, will have a finite onset temperature. At lower temperatures, the conductivity may be metallic or insulating, depending on t . In the former case, we have a ferromagnetic metal. In the latter case, the system becomes insulating at lower temperatures. It is possible that the ferromagnetic order is destroyed by the onset of insulating behavior. This cannot be resolved by the methods in this paper; thus, we have left the phase boundary of the paramagnetic insulating state intentionally vague. (In fact, it may not be paramagnetic, but might instead be a spin glass or some other non-ferromagnetic phase.)




FIG. 4: Phase diagram in $2 + \epsilon$ dimensions. PM is the paramagnetic metal phase; FM, the ferromagnetic metal phase; FI, the ferromagnetic insulator. PI is a non-ferromagnetic insulator: possibly paramagnetic, but perhaps a spin glass.

X. DISCUSSION

In this paper, we have used a straightforward and absolutely standard technique – albeit in a somewhat complicated and technically-involved context – to argue that

the instability of the weakly-interacting electron system in a weak random potential signals the development of ferromagnetism. Since there is a single instability occurring in the weak-interaction limit and it occurs while the resistivity is small, we are afforded a great deal of control over this problem. The instability bespeaks of the relative importance of one term in the action compared to the others. We need only satisfy this term – which we do by allowing for the development of a ferromagnetic moment – without worrying about other terms, since none of them are relevant. The demonstrable absence of a competing instability is the main justification for a simple mean-field theory. This may be formalized in a large- N generalization of the problem. Further credence is given by the physical arguments of the introductory section which rule out any other resolution of the divergence associated with the triplet amplitude. This is, in fact, a remarkable state of affairs, given that there is no comparable controlled theory for itinerant electron ferromagnetism in clean metals. In a clean metal, the Stoner instability occurs at strong coupling, where there are other competing instabilities and no unambiguous way of deciding between them.

It is striking how little is known about the phase diagram of the 2DEG. Even at zero magnetic field, the phase diagram as a function of r_s and disorder strength is only known in the small and large r_s limits for vanishing disorder. Intermediate r_s is poorly understood, as is finite disorder. We believe that we have, in this paper, extended *terra cognita* to include the small r_s , weak disorder region.

We emphasize that our result does not shed too much light on the metal-insulator transition which is observed in strongly-interacting 2DEGs. The development of ferromagnetism does not lead to a metallic state in two dimensions. Once ferromagnetism has developed, weak localization (which we have neglected in this paper) and interaction corrections both drive the system insulating. There is no contradiction with experiments, however, since the transition seen there is in the strong-interaction, high-resistivity ($t \sim 1$) regime where our result is not expected to be valid. On the other hand, our result *does* rule out theories which conjecture that the divergence of the triplet interaction leads to a metallic state *at zero-temperature* in 2D.

Acknowledgments

We would like to thank C. Chamon, A. Kamenev, E. Mucciolo, L. Radzihovsky, R. Shankar, and T. Senthil for discussions. This work was supported by the National Science Foundation under Grant No. DMR-9983544 and by the A.P. Sloan Foundation.

APPENDIX A: CORRESPONDENCE BETWEEN RG EQUATIONS AND AN EFFECTIVE ACTION

When we transform an action by introducing Hubbard-Stratonovich fields and integrating out the original variables of the action, the result can be an action which looks very different from the original one. However, the coupling constants of the two actions are related, and the corresponding RG equations should be as well. In the simplest cases, this is completely transparent. Consider, for instance, a system of N flavors of spinless fermions in one-dimension. Suppose that their density is commensurate with the underlying crystal lattice and that their interaction respects the $U(N)$ symmetry which mixes the fermions:

$$\mathcal{L} = \psi_{aR}^\dagger i(\partial_t + \partial_x) \psi_{aR} + \psi_{aL}^\dagger i(\partial_t - \partial_x) \psi_{aL} + \frac{g}{N} \left(\psi_{aR}^\dagger \psi_{aL} + \psi_{aL}^\dagger \psi_{aR} \right)^2 \quad (\text{A1})$$

Here, $a = 1, 2, \dots, N$. ψ_{aR}, ψ_{aL} annihilate fermions of flavor a at, respectively, the right and left Fermi points. This is the $U(N)$ Gross-Neveu model. Let us consider its large- N limit.

First, consider the RG equation for g . There is a single one-loop diagram, depicted on the left of fig. 5. All other diagrams are smaller by powers of $1/N$. The resulting RG equation is:

$$\frac{dg}{d\ell} = \frac{g^2}{2\pi} \quad (\text{A2})$$




FIG. 5: One-loop diagrams contributing to the RG equation for g .

If we introduce a Hubbard-Stratonovich field Δ which decouples the interaction

$$\mathcal{L} = \psi_{aR}^\dagger i(\partial_t + \partial_x) \psi_{aR} + \psi_{aL}^\dagger i(\partial_t - \partial_x) \psi_{aL} + \Delta \left(\psi_{aR}^\dagger \psi_{aL} + \psi_{aL}^\dagger \psi_{aR} \right) + \frac{N}{2g} \Delta^2 \quad (\text{A3})$$

then we can integrate out ψ :

$$\mathcal{L}_{\text{eff}} = \frac{N}{2g} \Delta^2 - N \text{tr} \ln ((i\partial_t - i\partial_x \sigma_3 + \Delta \sigma_1) \delta_{ab}) \quad (\text{A4})$$

where we have combined ψ_{aR}, ψ_{aL} into a two-component spinor on which $\sigma_{1,3}$ act. Δ cuts off the infrared logarithmic divergence of the $\text{tr} \ln$ term.

$$\mathcal{L}_{\text{eff}} = \frac{N}{2g} \Delta^2 - \frac{1}{4\pi} N \Delta^2 (\ln(\Lambda/\Delta) - 1)$$

$$= \left(\frac{N}{2g} - \frac{1}{4\pi} \ln(\Lambda/\Delta) + \frac{1}{4\pi} \right) \Delta^2 \quad (\text{A5})$$

where Λ is the ultraviolet cutoff. The second line suggests that $\frac{1}{4\pi} \ln(\Lambda/\Delta)$ renormalizes $1/g$. Thus, the effective action manifestly embodies the physics contained in the RG equation (A2). This is also clear from the close kinship between the one-loop diagram which gives the $\text{tr} \ln(\dots)$ in (A4) and the one-loop diagram which yields the RG equation (A2).

However, strictly speaking, the formal relationship between the RG equation and the effective action (A5) follows from the cutoff-independence of physically-measurable quantities, Δ being the simplest. From (A5), we see that the saddle-point value of Δ is

$$\Delta = \Lambda e^{-2\pi/g} \quad (\text{A6})$$

Differentiating both sides with respect to Λ and requiring that $d\Delta/d\Lambda$ vanish, we find:

$$\Lambda \frac{dg}{d\Lambda} = -\frac{g^2}{2\pi} \quad (\text{A7})$$

which is equivalent to the RG equation (A2).

Now, consider a slightly more complicated situation:

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_0(\psi, \psi^\dagger) + \mathcal{L}_0(\chi, \chi^\dagger) + \mathcal{L}_{\text{int}}(\chi, \chi^\dagger) \\ & + \frac{g}{N} \left(\psi_{aR}^\dagger \psi_{aL} + \psi_{aL}^\dagger \psi_{aR} \right)^2 \\ & + \frac{v}{\sqrt{N}} \left(\psi_{aR}^\dagger \psi_{aL} + \psi_{aL}^\dagger \psi_{aR} \right) \left(\chi_{aR}^\dagger \chi_{aL} + \chi_{aL}^\dagger \chi_{aR} \right) \end{aligned} \quad (\text{A8})$$

where $\mathcal{L}_0(\psi, \psi^\dagger)$ is the same as the first line of (A1).

The RG equation for g now reads:

$$\frac{dg}{d\ell} = \frac{g^2}{2\pi} + \frac{v^2}{2\pi} \quad (\text{A9})$$

The second term arises from the diagram on the right-hand-side of fig. 5.

Now, consider $\mathcal{L}_{\text{eff}}(\Delta, \chi, \chi^\dagger)$, where Δ is introduced to decouple the g interaction, as before. We can no longer calculate it exactly. However, if we assume that $v \ll g$, we can calculate it to lowest non-trivial order in v . The lowest order diagram by which it contributes is the second one of fig. 6. We integrate out ψ completely and integrate out all χ fluctuations above some cutoff Λ' . We are free to choose Λ' as we like, of course, but the natural choice is $\Lambda' = \Delta$ so that our effective action is for excitations with energies of order Δ . The resulting effective action is

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & \frac{N}{2g} \Delta^2 - \frac{1}{4\pi} N \Delta^2 \ln(\Lambda/\Delta) \\ & - \frac{1}{4\pi} v^2 N \Delta^2 (\ln(\Lambda/\Delta))^3 \\ & + \mathcal{L}_0(\chi, \chi^\dagger) + \mathcal{L}_{\text{int}}(\chi, \chi^\dagger) \end{aligned} \quad (\text{A10})$$

The last term looks strange. It looks like an $O(v^2)$ renormalization of $1/g$, which would imply a contribution of order $g^2 v^2$ to $dg/d\ell$. Also, this term is proportional to $(\Delta^2 \ln(\Lambda/\Delta))^3$ rather than $\Delta^2 \ln(\Lambda/\Delta)$ as we might have expected from the second diagram of fig. 5 and the RG equation (A9).




FIG. 6: The one- and three-loop contributions to the effective action for Δ which encapsulate the same physics as the one-loop RG equation.

The problem is that we are looking at $\mathcal{L}_{\text{eff}}(\Delta)$, which is not observable, rather than the vacuum expectation value of Δ , which is. When we compute Δ , the two difficulties

noted above resolve each other. Minimizing the effective action, we find that, to lowest non-trivial order in v :

$$\ln(\Lambda/\Delta) = \frac{2\pi}{g} - \frac{2\pi v^2}{g^3} \quad (\text{A11})$$

or

$$\Delta = \Lambda e^{-2\pi/(g+v^2/g)} \quad (\text{A12})$$

Requiring Λ -independence of Δ , we recover the RG equation for g – and also the RG equation for v which we didn't compute.

The lesson is that a one-loop contribution to the RG equation for a coupling constant might manifest itself in an effective action only through a higher-loop (in the above example, three-loop) diagram. In the interacting electron problem in a random potential, diagrams (a), (b), and (c) could enter $S_{\text{eff}}[X]$ through three-loop diagrams.

-
- ¹ B. L. Altshuler and A. G. Aronov, Solid State Comm. **39**, 115 (1979); Sov. Phys. JETP **50**, 968 (1979).
 - ² A. M. Finkelstein, Zh. Eksp. Teor. Fiz. **84**, 168 (1983), Sov. Phys. JETP **57**, 97 (1983).
 - ³ C. Castellani, C. Di Castro, P. A. Lee and M. Ma, Phys. Rev. B **30** 527 (1984).
 - ⁴ C. Castellani, C. Di Castro, P. A. Lee, M. Ma, S. Sorella, and E. Tabet, Phys. Rev. B **30**, 1596 (1984).
 - ⁵ D. Belitz and T. R. Kirkpatrick, Rev. Mod. Phys. **66**, 261 (1994).
 - ⁶ A. M. M. Pruisken, M. A. Baranov, and B. Škorić, cond-mat/9712322, 9712323.
 - ⁷ C. Chamon, A. W. W. Ludwig, and C. Nayak, Phys. Rev. B **60**, 2239 (1998).
 - ⁸ Note that Finkelstein uses g to denote resistance, as did Chamon, Ludwig, and Nayak, in keeping with his notation. However, the more widespread usage is for g to denote conductance. To avoid confusion, we adopt the notation of Castellani, Di Castro, Lee, and Ma: t denotes resistance.
 - ⁹ S. V. Kravchenko *et al.*, Phys. Rev. B **51**, 7038 (1995); D. V. Kravchenko *et al.*, Phys. Rev. Lett. **77**, 4938 (1996); D. Simonian, S. V. Kravchenko, M. P. Sarachik, and V. M. Pudalov, Phys. Rev. Lett. **79**, 2304 (1997).
 - ¹⁰ D. Popović, A. B. Fowler, and S. Washburn, Phys. Rev. Lett. **79**, 1543 (1997).
 - ¹¹ Y. Hanien *et al.*, Phys. Rev. Lett. **80**, 1288 (1998); M. Y. Simmons *et al.*, *ibid*, **80**, 1292 (1998); S. J. Papadakis and M. Shayegan, Phys. Rev. B, June 15 (1998).
 - ¹² J. Lam *et al.*, Phys. Rev. B **56**, R12741 (1997); P. T. Coleridge *et al.*, *ibid*, **56**, R12764 (1997).
 - ¹³ P. Phillips, Y. Wan, I. Martin, S. Knysh, and D. Dalidovich, Nature **395**, 253 (1998); P. Phillips, cond-mat/9910122.
 - ¹⁴ Q. Si and C. Varma, Phys. Rev. Lett. **81**, 4951 (1998).
 - ¹⁵ S. Chakravarty, S. Kivelson, C. Nayak, and K. Voelker, Phil. Mag. B **79** 859 (1999).
 - ¹⁶ B. Altshuler and D. Maslov, Phys. Rev. Lett. **82**, 145 (1999).
 - ¹⁷ S. Das Sarma and E. H. Hwang, Phys. Rev. Lett. **83**, 164 (1999); *ibid* **84**, 5596 (2000).
 - ¹⁸ C. Chamon, E. R. Mucciolo, A. H. C. Neto, Phys. Rev. B **64**, 245115/1-12 (2001).
 - ¹⁹ When the bare interaction is weak, disorder does not allow the former, an SDW, and its necessary concomitant, a CDW, to form since the energy scale at which the interaction becomes strong is small compared to the inverse of the elastic scattering time. On the other hand, one can imagine that strong interactions could lead to the formation of an SDW in the ballistic regime, i.e. at length scales shorter than the mean-free path. At longer scales, SDW order would be disrupted by impurities, but the correlation length could be long if the impurity density is low. In the problem which we consider, however, interactions are assumed to be weak in the ballistic regime. They only become strong in the diffusive regime, where SDW order will not develop.
 - ²⁰ R. N. Bhatt and P. A. Lee, Phys. Rev. Lett. **48**, 344 (1982).
 - ²¹ What we really mean by this is that spin glass order cannot be important *at this scale*. However, local ferromagnetism, which develops at the scale at which the triplet interaction becomes strong, could be disrupted at longer scales (at which the effective disorder strength is large) so that spin glass order – e.g. a cluster spin glass – forms at the longest length scales. Furthermore, in an insulating state, caused by Wigner crystallization and/or Anderson localization, a spin glass could form since there will be competition between different multi-spin exchange process which lead to effective spin-spin interactions of different signs.
 - ²² B. N. Narozhny, I. L. Aleiner, and A. I. Larkin, Phys. Rev. B **62**, 14898 (2000).
 - ²³ A. V. Andreev and A. Kamenev, Phys. Rev. Lett. **81**, 3199 (1998).
 - ²⁴ C. Chamon and E. Mucciolo, Phys. Rev. Lett. **85**, 5607 (2000).

- ²⁵ D. Belitz and T. R. Kirkpatrick, Phys. Rev. B **53**, 14364 (1996).
- ²⁶ O. Prus, Y. Yaish, M. Reznikov, U. Sivan, and V. Pudalov, cond-mat/0209142.
- ²⁷ N. Bonesteel, I. A. McDonald, and C. Nayak, Phys. Rev. Lett. **77**, 3009 (1996).
- ²⁸ N.Read and S.Sachdev, Phys. Rev. Lett. **75**, 3509 (1995).